



Vlaams Instituut voor de Zee
Flanders Marine Institute

日本及び印度太平洋産貝類目録(2)

腹足綱(ソデガイ超科)

波部忠重・小菅貞男

34929

A List of the Indo-Pacific Molluscs,
concerning to the Japanese Molluscan Fauna (2)

Class Gastropoda (Superfamily Stromboidea)

by

Tadashige HABE and Sadao KOSUGE

国立科学博物館
東京, 上野公園
(振替東京 55629 : 波部忠重)

c/o National Science Museum
Ueno Park, Tokyo, Japan
June, 1964

Superfamily Stromboidea ソデガイ超科

Family Strombidae ソデガイ科

Genus *Tibia* [Röding 1798] ハシナガソデガイ属

Syn. *Rostellaria* Lamarck 1799, *Gladius* H. & A. Adams 1854.

Subgenus *Varicospira* Eames 1952 カゴメソデガイ亜属

1. *T. (V.) cancellata* (Lamarck, 1822) (*Strombus*) カゴメソデガイ
Syn. *Strombus crista* Sowerby, 1842; *Rimella speciosa* H. & A. Adams, 1863; *R. tayleri* H. & A. Adams, 1863; *R. cancellata minor* Schepman, 1909.
Loc.: Indonesia; Philippines; China Sea.

Subgenus *Tibia* (s. s.) ハシナガソデガイ亜属

2. *T. (T.) curta* (Sowerby, 1822) (*Rostellaria*). ダイコクボラ
Loc.: Persian Gulf; India.
3. *T. (T.) insulaechorab* [Röding, 1798] エビスボラ
Syn. *Rostellaria curvirostris* Lamarck, 1822; *R. brevisrostra* Schumacher, 1817; *R. magna* Schröter, 1788 (n. n.); *R. ternatarium* Montfort, 1810; *R. dentata* Perry, 1811; *R. luteostoma* Angas, 1878.
Loc.: Red Sea; Molucca; India.
4. *T. (T.) fusus* (Linné, 1767) (*Strombus*) ハシナガソデガイ
Syn. *Strombus clavus* Gmelin, 1817; *St. unicornis* Dillwyn, 1817; *Rostellaria subulata* Schumacher, 1817; *R. rectirostris* Lamarck, 1822; *T. indiarum* Röding, 1798; *T. contorta* Röding, 1798.
Loc.: Philippines; Formosa; China.
5. *T. (T.) melanochilus* (A. Adams, 1855) (*Gladius*) クチグロハシナガソデガイ
Loc.: China.
6. *T. (T.) powisii* (Petit, 1842) (*Rostellaria*) ヤマビトボラ
Syn. *Rostellaria powisii abyssicola* Schepman, 1909.
Loc.: Indonesia; Philippines; Formosa; Kyushu; Shikoku; Honshu (north to Kii Peninsula). 紀伊半島以南.
7. *T. (T.) serrata* (Perry, 1811) (*Rostellaria*) ノコギリボラ
Syn. *Strombus fissus* Dillwyn, 1817; *Rostellaria favanni* Pfeiffer, 1840.
Loc.:? (不明).
8. *T. (T.) martinii* (Marrat, 1877) (*Gladius*) ワタナベボラ
Loc.: Philippines.
Subgenus *Rostellariella* Thiele 1929 ベンガルソデガイ亜属
9. *T. (R.) delicatula* (Nevill, 1881) (*Rostellaria*) ベンガルソデガイ
Loc.: Persian Gulf; Bay of Bengal.

Genus *Terebellum* [Röding 1798] トンボガイ属

10. *T. terebellum* (Linné, 1767) (*Conus*)

a. *T. terebellum terebellum* (Linné, 1767) トンボガイ

Syn. *T. lineatum* Röding, 1798; *T. nebulosum* Röding, 1798;

T. punctolorum Röding, 1798; *T. subulatum* Lamarck, 1822; *T. variegatum* Link, 1807; *T. maculosum* A. Adams, 1850.

Loc.: North Australia; New Caledonia; New Guinea; Philippines; Ryukyu; Amami Islands. 奄美以南.

b. *T. terebellum delicatum* Kuroda & Kawamoto, 1956 ウストンボガイ

Loc.: Kyushu; Shikoku; Honshu (North to Bôshû Peninsula). 房州以南.

Genus *Strombus* Linné 1758 ソデガイ属

Syn. *Strombella* Schlüter 1838.

Subgenus *Strombus* Linné ソデガイ亜属

*11. *St. (St.) pugilis* Linné, 1758 ソデガイ

a. *St. (St.) pugilis pugilis* Linné, 1758 ソデガイ

Syn. *Strombus pugilis peculiaris* Smith, 1940.

Loc.: Florida; West Indies; Central America; South America (South to Brazil).

b. *St. (St.) pugilis alatus* Gmelin, 1790 ソデガイモドキ

Loc.: South eastern coast of North America.

c. *St. (St.) pugilis nicaraguensis* Fluck, 1905

Loc.: Nicaragua.

*12. *St. (St.) gracilior* Sowerby, 1825 ニシノソデボラ

Loc.: Gulf of California to Peru (South America).

Subgenus *Laevistrombus* Kira 1955 スイショウガイ亜属

13. *St. (L.) canarium* Linné, 1758

a. *St. (L.) canarium canarium* Linné, 1758. サザナミスイショウガイ

Syn. *St. vanikorensis* Quoy & Gaimard, 1834.

Loc.: New Caledonia; Solomon Is.; North-east Australia; New Guinea; Borneo; Indonesia; Philippines; Formosa; Ryukyu; Amami Islands; Ceylon. 奄美以南.

b. *St. (L.) canarium turturella* [Röding, 1798] (*Lambis*) スイショウガイ

Syn. *Strombus isabella* Lamarck, 1822; *St. taeniatus* Quoy & Gaimard, 1834.

Loc.: Indonesia; Philippines; Ryukyu; Kyushu; Shikoku; Honshu (north to Sagami Bay). 相模湾以南.

Subgenus *Tricornis* Jousseaume 1886 ゴホウラ亜属

Syn. *Lobatus* Iredale 1921, *Aliger* Thiele 1929, *Eustrombus* Wenz 1940.

14. *St. (T.) tricornis* Humphrey, 1786 ミツカドソデガイ
Syn. *Strombus orientalis* Duclos, 1844; *St. pertinax* Duclos, 1844.
Loc.: Red Sea; Aden (East coast of Africa).
15. *St. (T.) thersites* Swainson, 1823 アツソデガイ
Syn. *Strombus ponderosus* Philippi, 1842.
Loc.: New Caledonia; Ryukyu; Amami Islands. 奄美以南.
16. *St. (T.) latissimus* Linné, 1758 ゴホウラ
Syn. *Lambis picta* Röding, 1798; *Pterocera alata* Schumacher, 1817.
Loc.: Fiji Is.; New Caledonia; Indonesia; Philippines; Formosa, Ryukyu;
Amami Is. 奄美以南.
17. *St. (T.) taurus* Reeve, 1857 コッテイソデガイ
Loc.: Marshall Is.; Mariana Is.
18. *St. (T.) sinuatus* Humphrey, 1786 ヒメゴホウラ
Syn. *Lambis lobata* Röding, 1798; *Pterocera palmata* Fischer, 1807;
Strombus laciniatus Dillwyn, 1817; *St. cristatus* Lamarck, 1822.
Loc.: Fiji Is.; New Caledonia; Solomon Is.; North-east Australia; New Guinea;
Celebes; Borneo; Marshall Is.; Caroline Is.; Philippines; Formosa; Ryukyu;
Amami Islands. 奄美以南.
- *19. *St. (T.) raninus* Gmelin, 1790 タカノハソデガイ
Syn. *Strombus bituberculata* Lamarck, 1822.
Loc.: Florida; West Indies; Colombia (South America).
- *20. *St. (T.) gallus* Linné, 1758 クタカケソデガイ
Loc.: West Indies; South America (South to Brazil).
- *21. *St. (T.) costatus* Gmelin, 1790 ミルクソデガイ
Syn. *Strombus accipitrinus* Lamarck, 1822; *St. inermis* Swainson, 1882;
St. samba Clench, 1937.
Loc.: Florida; West Indies; Central America; South America (south to Brazil).
- *22. *St. (T.) gigas* Linné, 1758 ピンクガイ
Syn. *Strombus gigas horridus* Smith, 1937; *S. canaliculatus* Burry, 1949.
Loc.: Bermuda; Southern Florida; West Indies; Northern South America.
- *23. *St. (T.) goliath* Sowerby, 1842 ダイオウソデガイ
Loc.: West Indies; West coast of South America.
- *24. *St. (T.) galeatus* Swainson, 1823 カブトソデガイ
Loc.: Gulf of California to Equador (South America).
- *25. *St. (T.) peruvianus* Swainson, 1823 ゴゼンソデガイ
Loc.: Mexico to Northern Peru (South America).

Subgenus *Canarium* Schumacher, 1817 オハグロガイ亜属

Syn. *Strombidea* Swainson 1840, *Conorium* Jousseume 1880.

26. *St. (C.) urceus* Linné, 1758

a. *St. (C.) urceus urceus* Linné, 1758 オハグロガイ

Syn. *Canarium ustulatum* Schumacher, 1817; *Strombus incisus* Wood, 1828; *St. anatellus* Duclos, 1844; *St. muricatus* Watson, 1885; *St. plicatus* Adams & Leloup, 1938 (non Lamarck, 1816); *St. ustulatus* form *laevis* Dodge, 1946.

Loc.: Fiji Is.; New Caledonia; Solomon Is.; North-east Australia; New Guinea; Celebes; Marshall Is.; Caroline Is.; Philippines; Formosa; Ryukyu; Amami Island; Honshu (north to Kii Peninsula). 紀伊半島以南.

b. *St. (C.) urceus orrae* Abbott, 1960 オドリソデガイ

Loc.: North-east Australia; Arafra Sea.

27. *St. (C.) labiatus* [Röding, 1798] (*Lambis*)

a. *St. (C.) labiatus labiatus* [Röding, 1798] フトスジムカシタモトガイ

Syn. *Lambis reticulata* Link, 1807; *Strombus plicatus* Lamarck, 1816; *St. dentatus* Reeve, 1850 (non Linné, 1758), *Canarium otiolium* Iredale, 1931; *Strombus ustulatus plicatus* Dodge, 1946 (non Lamarck, 1816); *Canarium dentatum* Kira, 1959 (non Linne, 1758).

Loc.: Fiji Is.; New Caledonia; New Hebrides; Solomon Is.; North-east Australia; New Guinea; Indonesia; Celebes; Philippines; Ryukyu; Amami Is.; Honshu (north to Kii Peninsula); Thailand; Andaman Is. 紀伊半島以南.

b. *St. (C.) labiatus olydius* Duclos, 1844

Loc.: Mauritius; Madagascar; Zanzibar; East coast of Africa; Ceylon.

28. *St. (C.) klineorum* Abbott, 1960

Syn. *Strombus urceus* Duclos, 1844 (non Linné, 1758)

Loc.: Ceylon.

29. *St. (C.) microurceus* (Kira, 1958) (*Canarium*) ヤサガタムカシタモトガイ

Loc.: Samoa Is.; Fiji Is.; Norfolk I.; Solomon Is.; North-east Australia; New Guinea; Bismark Is.; Indonesia; Caroline Is.; Mariana Is.; Philippines; Ryukyu; Amami Is.; Kyushu; Shikoku; Honshu (north to Kii Peninsula). 紀伊半島以南.

30. *St. (C.) mutabilis* Swainson, 1821

a. *St. (C.) mutabilis mutabilis* Swainson, 1821 ムカシタモトガイ

Syn. *Strombus floridus* Lamarck, 1822; *St. epimellus* Duclos, 1844; *St. flosculosus* Mörch, 1852; *St. flammeus* Issel & T-Canefri, 1876; *St. floridus zebriolata* Adams & Leloup, 1938.

Loc.: Samoa Is.; North Australia; Indonesia; New Caledonia; Fiji Is.; New

- Hebrides; Solomon Is.; New Guinea; Mariana Is.; Caroline Is.; Philippines; Ryukyu; Amami Is.; Hachijo I.; Honshu (North to Kii Peninsula); Andaman Is.; Maldive Is.; Red Sea; Madagascar; East coast of Africa. 紀伊半島以南.
- b. *St. (C.) mutabilis ochroglottis* Abbott, 1960
 Loc.: Mauritius.
31. *St. (C.) maculatus* Sowerby, 1842 シロムカシタモトガイ
 Syn. *Strombus floridus depauperata* Dautzenberg & Bouge, 1933.
 Loc.: Easter I.; Tahite I.; Gilbert Is.; Marshall Is.; Caroline Is.; Hawaiian Is.
32. *St. (C.) fusiformis* Sowerby, 1842
 Loc.: Red Sea; Seychelles Is.; Madagascar; East coast of Africa.
33. *St. (C.) erythrinus* Dillwyn, 1817
 a. *St. (C.) erythrinus erythrinus* Dillwyn, 1817 チリメンムカシタモトガイ
 Syn. *Strombus elegans* Sowerby, 1842; *St. radians* Duclos, 1844; *St. ruppelli* Reeve, 1850; *Rostellaria rubicunda* Mathews & Iredale, 1912 (non Perry, 1810); *Strombus plicatus* Oostingh, 1925 (non Lamarck, 1816); *St. rugosus* Dietrich & Morris, 1953 (non Sowerby, 1825)
 Loc.: New Caledonia; Hawaiian Is.; Gilbert Is.; Marshall Is.; New Guinea; Indonesia; Caroline Is.; Mariana Is.; Philippines; Formosa; Ryukyu; Amami Is.; Red Sea; Madagascar; Réunion; Zanzibar (East coast of Africa). 奄美以南.
- b. *St. (C.) erythrinus rugosus* Sowerby, 1825
 Syn. *Strombus jugosus* Wood, 1828; *St. corrugatus* Adams & Reeve, 1848.
 Loc.: Ellice Is.; Samoa Is.; Fiji Is.; Tonga Is.
34. *St. (C.) haemastoma* Sowerby, 1842
 Loc.: Cook Is.; Samoa I.; Ellice Is.; Marshall Is.; Réunion; Mauritius.
35. *St. (C.) scalariformis* Duclos, 1833
 Loc.: Philippines; South China Sea.
36. *St. (C.) helli* Kiener, 1843
 Syn. *Strombus cancellatus* Pease, 1860.
 Loc.: Hawaiian Is.
37. *St. (C.) dentatus* Linné, 1758 ミツユビガイ
 Syn. *Strombus tridentatus* Gmelin, 1771; *Lambis dentata* Röding, 1798; *Strombus samar* Dillwyn, 1817; *St. samarensis* Reeve, 1851.
 Loc.: Society Is.; New Caledonia; Solomon Is.; Marshall Is.; Caroline Is.; New Guinea; Philippines; Formosa; Ryukyu; Amami Is.; Kyushu; Shikoku; Honshu (north to Kii Peninsula); Réunion; East coast of Africa. 紀伊半島以南.

38. *St. (C.) fragilis* [Röding, 1798] (*Lambis*) マクラソデガイ
 Syn. *Strombus samar* Dillwyn, 1817 (part); *St. dubius* Swainson, 1828; *St. bulbulus* Sowerby, 1842; *St. terebellatus* Abrard, 1946 (non Sowerby, 1842).
 Loc.: Samoa Is.; Fiji Is.; New Hebrides; Marshall Is.; Hawaiian Is.; Caroline Is.; Mariana Is.; New Guinea; Philippines; Ryukyu; Amami Is. 奄美以南.
39. *St. (C.) terebellatus* Sowerby, 1842
 a. *St. (C.) terebellatus terebellatus* Sowerby, 1842 トンボンソデガイ
 Syn. *Strombus dentatus* Kiener, 1843 (non Linné, 1758)
 Loc.: Fiji Is.; New Hebrides; Ellice Is.; Solomon Is.; New Guinea; Caroline Is.; Philippines; Ryukyu; Amami Is. 奄美以南.
 b. *St. (C.) terebellatus afrobellatus* Abbott, 1942
 Loc.: Red Sea; Madagascar; East coast of Africa.

Subgenus *Dolomena* Iredale 1931 オハグロシドロ亜属

40. *St. (D.) plicatus* [Röding, 1798] (*Lambis*)
 a. *St. (D.) plicatus plicatus* [Röding, 1798] アツシドロガイ
 Syn. *Strombus deformis* Griffith & Pidgeon, 1834; *St. sibbaldi* Kiener, 1834 (non Sowerby, 1842).
 Loc.: Red Sea.
 b. *St. (D.) plicatus columba* Lamarck, 1822
 Syn. *Strombus tankervillei* Swainson, 1823.
 Loc.: Seychelles Is.; Madagascar; East coast of Africa.
 c. *St. (D.) plicatus sibbaldi* Sowerby, 1842
 Syn. *Strombus deformis* Kiener, 1843 (non Gray, 1834); *St. kieneri* Issel & T-Canefri, 1876; *St. yerburyi* Smith, 1891.
 Loc.: Ceylon; Aden; Arabia.
 d. *St. (D.) plicatus pulchellus* Reeve, 1851 オハグロシドロガイ
 Syn. *Strombus malekulensis* Abrard, 1946.
 Loc.: Fiji Is.; New Hebrides; Solomon Is.; New Guinea; Caroline Is.; Philippines; Ryukyu; Amami Is.; Kyushu; Shikoku; Honshu (North to Ise Bay). 伊勢湾以南.
41. *St. (D.) dilatatus* Swainson, 1821
 a. *St. (D.) dilatatus dilatatus* Swainson, 1821 カネツケシドロガイ
 Syn. *Strombus orosminus* Duclos, 1844.
 Loc.: New Caledonia; New Hebrides; Solomon Is.; New Guinea; Malaya; Philippines; Ryukyu; Shikoku. 四国以南.
 b. *St. (D.) dilatatus swainsoni* Reeve, 1850
 Loc.: ? (不明).

42. *St. (D.) labiosus* Wood, 1828 コブシドロガイ
 Loc.: Indonesia; Borneo; New Guinea; Philippines; Maldive Is.; Madagascar;
 East coast of Africa.
43. *St. (D.) marginatus* Linné, 1758
 a. *St. (D.) marginatus marginatus* Linné, 1758
 Syn. *Lambis carinata* Röding, 1798.
 Loc.: Sumatra (Indonesia); Burma; Ceylon.
 b. *St. (D.) marginatus succinctus* Linné, 1767
 Syn. *Strombus accinctus* Linné, 1767.
 Loc.: Ceylon; Madras (Southern India).
 c. *St. (D.) marginatus robustus* Sowerby, 1874 フドロガイ
 Syn. *Strombus septimus* Duclos, 1899; *Laevistrombus succinctus* Kira, 1959
 (non Linné, 1767).
 Loc.: Indochina; Kyushu; Shikoku; Honshu (north to Bôsô Peninsula);
 Malaya; Burma. 房総以南.
 d. *St. (D.) marginatus septimus* Duclos, 1844 ヨロイノソデガイ
 Loc.: New Caledonia; Solomon Is.; New Guinea; Philippines; Ryukyu;
 Amami Is. 奄美以南.
44. *St. (D.) variabilis* Swainson, 1820 ホクロソデガイ
 a. *St. (D.) variabilis variabilis* Swainson, 1820 ホクロソデガイ
 Syn. *Strombus lituratus* Menke, 1829.
 Loc.: North-east Australia, Celebes; Indonesia; Malaya; Philippines; Ryukyu;
 Amami Is. 奄美以南.
 b. *St. (D.) variabilis athenius* Duclos, 1844
 Syn. *Strombus variabilis* Kiener, 1843 (non Swainson)
 Loc.: New Caledonia; Fiji Is.; Samoa Is.; Marshall Is.; Solomon Is.;
 New Guinea; Caroline Is.
45. *St. (D.) minimus* Linné, 1771 コソデガイ
 Syn. *Lambis minimus* Röding, 1798; *Strombus troglodytes* Lamarck, 1822.
 Loc.: New Caledonia; Fiji Is.; New Hebrides; Solomon Is.; New Guinea;
 Indonesia; Philippines; Formosa; Ryukyu; Amami Is. 奄美以南.

Subgenus *Labiostrombus* Oostingh 1925. ホカケソデガイ亜属

Syn. *Gallinula* Herrmannsen 1847.

46. *St. (L.) epidromis* Linné, 1758 ホカケソデガイ
Syn. *Lambis epidromis* Röding, 1798; *Strombus expansa* Tryon, 1885.
Loc.: New Caledonia; North-east Australia; New Guinea; Malaya; Philippines;
Ryukyu; Amami Is. 奄美以南.

Subgenus *Doxander* Iredale 1931 シドロガイ亜属

47. *St. (D.) vittatus* Linné, 1758
a. *St. (D.) vittatus vittatus* Linné, 1758 タケノコシドロガイ
Syn. *Lambis vittatus* Röding, 1798; *Strombus australis* Schröter, 1805;
St. turritus Lamarck, 1822.
Loc.: Fiji Is.; New Hebrides; New Guinea; Celebes; Borneo; Malaya; Phi-
lippines; Hong Kong; Ryukyu. 沖縄以南.
b. *St. (D.) vittatus japonicus* Reeve, 1851 シドロガイ
Syn. *Lora suturolineata* Nomura & Hatai, 1940.
Loc.: Formosa; Kyushu; Shikoku; Honshu (North to Mutsu Bay). 陸奥湾以南.
c. *St. (D.) vittatus campbelli* Griffith & Pidgeon, 1834
Syn. *Alaba (Styliferina) sulcata* Watson, 1886.
Loc.: North-east Australia; Arafra Sea; North-west Australia.
48. *St. (D.) listeri* Gray, 1852 ヒメゴゼンガイ
Syn. *Strombus mirabilis* Sowerby, 1870.
Loc.: Ceylon; Andaman Is.

Subgenus *Lentigo* Jousseume 1886 イボソデガイ亜属

49. *St. (L.) lentiginosus* Linné, 1758 イボソデガイ
Syn. *Lambis rana* Röding, 1798.
Loc.: Society Is.; Samoa Is.; Fiji Is.; New Caledonia; New Hebrides; Marshall
Is.; Solomon Is.; North-east Australia; New Guinea; Celebes; Caroline Is.;
Philippines; Formosa; Ryukyu; Amami Is.; Ceylon; Malaya; Cocos Is.;
Nicobar Is.; Maldive Is.; Seychelles Is.; Mauritius; Réunion; Madagascar;
East coast of Africa. 奄美以南.
50. *St. (L.) pipus* [Röding, 1798] (*Lambis*) オハグロイボソデガイ
Syn. *Strombus papilio* Dillwyn, 1817; *St. exustus* Swainson, 1822; *St.*
adustus Reeve, 1851.
Loc.: Society Is.; Solomon Is.; Queensland, Australia; Band & Timor Is,
Indonesia; Philippines; Formosa; Ryukyu; Seychelles Is.; Réunion; Zanzibar
(East coast of Africa). 沖縄以南.

51. *St. (L.) fasciatus* Born, 1778 ハチマキノデガイ
 Syn. *Strombus subalata* Herbst, 1778; *Lambis elegantissima* Röding, 1798;
St. polyfasciatus Dillwyn, 1817; *St. lineatus* Lamarck, 1822; *St. lineolatus*
 Wood, 1828; *St. flavigula* Tryon, 1885 (non Meuschen).
 Loc.: Red Sea.
- *52. *St. (L.) latus* Gmelin, 1791 ヒロソデガイ
 Loc.: West Africa.
- *53. *St. (L.) granulatus* Swainson, 1828 メキシコソデガイ
 Loc.: Gulf of California to Equador (South America) .

Subgenus *Euprotomus* Gill 1870 マイノソデガイ亜属

54. *St. (E.) aurisdianae* Linné, 1758
- a. *St. (E.) aurisdianae aurisdianae* Linné, 1758 マイノソデガイ
 Syn. *Lambis stiva* Röding, 1798; *L. buris* Röding, 1798; *Strombus lamarckii*
 Sowerby, 1842 (non Gray, 1826); *St. striatogranosus* Martens, 1880; *St.*
aurisdianae chrysostomus Kuroda, 1942 (ヤヨイソデガイ).
 Loc.: Solomon Is.; New Guinea; Celebes; Indonesia; Malaya; Philippines;
 Formosa; Ryukyu; Amami Is.; Ceylon; Seychelles Is.; Réunion. 奄美以南.
- b. *St. (E.) aurisdianae aratrum* [Röding, 1798] (*Lambis*) クリフマイノソデガイ
 Syn. *Strombus aurisdianae* Lamarck, 1816 (non Linné); *St. aurisdianae*
adusta Dillwyn, 1817; *St. melastomus* Swainson, 1822; *St. aurisasimi* Dillwyn,
 1823; *St. melanostomus* Sowerby, 1825; *St. adusta* Gray, 1826.
 Loc.: North-east Queensland, Australia.
55. *St. (E.) bulla* [Röding, 1798] (*Lambis*) ベニソデガイ
 Syn. *Strombus laevis* Perry, 1811; *St. lamarckii* Gray, 1826 (non Sowerby,
 1842); *St. aurisdianae* Reeve, 1842 (non Linné, 1758); *St. guttatus* Kiener,
 1843; *St. aurisdianae* Oostingh, 1925 (non Linné).
 Loc.: Samoa Is.; New Caledonia; Solomon Is.; New Guinea; Celebes; Indonesia;
 Philippines; Formosa; Ryukyu; Amami Is. 奄美以南.
56. *St. (E.) vomer* [Röding, 1798] (*Lambis*)
- a. *St. (E.) vomer vomer* [Röding, 1798] ウラスジマイノソデガイ
 Syn. *Strombus acutus* Perry, 1811; *St. pacificus* Swainson, 1821; *St.*
zelandiae Gray, 1826; *St. chemnitzii* Pfeiffer, 1840; *St. novaezelandiae*
 Reeve, 1842; *St. novaezelandia* Paetel, 1869; *St. atratum* Allan, 1950;
St. vomer hirasei Kuroda, 1942 (ヒラセソデガイ).
 Loc.: New Caledonia; Philippines; Ryukyu; Amami Is. 奄美以南.
- b. *St. (E.) vomer hawaiiensis* Pilsbry, 1917
 Loc.: Hawaiian Is.

- c. *St. (E.) vomer iredalei* Abbott, 1960
 Syn. *Strombus australis* Gray, 1826 (non Schröter); *Strombus australis* Sowerby, 1842 (non Schröter).
 Loc.: North-west Australia.

Subgenus *Conomurex* Fischer 1884 マガキガイ亜属

57. *St. (C.) luhuanus* Linné, 1758 マガキガイ
 Syn. *Strombus luhuanus* Herbst, 1788; *Lambis luhuana* Röding, 1798
 Loc.: New Caledonia; Fiji Is.; Queensland, Australia; Solomon Is.; New Guinea; Marshall Is.; Caroline Is.; Celebes; Malaya; Philippines; Formosa; Ryukyu; Kyushu; Shikoku; Honshu (north to Bôso Peninsula). 房総以南.
58. *St. (C.) decorus* [Röding, 1798] (*Lambis*)
 a. *St. (C.) decorus decorus* [Röding, 1798]
 Syn. *Lambis miniata* Link, 1807; *L. flammea* Link, 1807; *Strombus cylindricus* Swainson, 1821; *St. mauritianus* Lamarck, 1822; *St. lutruanus* Dillwyn, 1823; *St. laevilabris* Menke, 1828; *St. coniformis* Sowerby, 1842.
 Loc.: Sumatra; Indonesia; Burma; Ceylon; Seychelles Is.; Mauritius; Réunion; Madagascar; East coast of Africa.
- b. *St. (C.) decorus persicus* Swainson, 1821 ペルシヤママガキガイ
 Syn. *St. ismarius* Duclos, 1844; *St. (Conomurex) belutschiensis* Melville, 1898.
 Loc.: Red Sea; Persian Gulf.

Subgenus *Gibberulus* Jousseau 1888 ネジマガキガイ亜属

59. *St. (G.) gibberulus* Linné, 1758
 a. *St. (G.) gibberulus gibberulus* Linné, 1758
 Syn. *Strombus labiatus* Perry, 1811.
 Loc.: Hong Kong (in error); Andaman Is.; Madras, India; Ceylon; Cocos Is.; Maldive Is.; Seychelles Is.; Mauritius; Madagascar; East coast of Africa.
- b. *St. (G.) gibberulus gibbosus* [Röding, 1798] (*Lambis*) ネジマガキガイ
 Syn. *Strombus gibberulus* Sowerby, 1842 (non Linné).
 Loc.: Society Is.; Cook Is.; Fiji Is.; New Caledonia; New Hebrides; Solomon Is.; Queensland, Australia; New Guinea; Celebes; Java; Marshall Is.; Caroline Is.; Mariana Is.; Philippines; Formosa; Ryukyu; Amami Is. 奄美以南.

- c. *St. (G.) gibberulus albus* Mörch, 1850
 Syn. *Strombus gibberulus rhodostomus* Martens, 1869.
 Loc.: Red Sea; Gulf of Aden.

Genus *Lambis* [Röding 1798] クモガイ属

Syn. *Pterocera* Lamarck 1799, *Pteroceras* Link 1807, *Pteroceres* Montfort 1810, *Digitata* Fabricius 1823, *Pterocerus* Brongniart 1829, *Heptadactylus* Mörch 1852.

Subgenus *Lambis* (S. S.) クモガイ亜属

60. *L. (L.) lambis* (Linné, 1758) (*Strombus*) クモガイ
 Syn. *Lambis lambis* Röding, 1798; *L. lamboides* Röding, 1798; *L. lobata* Röding, 1798; *L. cerea* Röding, 1798; *L. maculata* Röding, 1798; *L. hermaphrodita* Röding, 1798; *L. laciniata* Röding, 1798; *Strombus camelus* Gray, 1826; *Pterocera crocata* Yonge, 1932 (non Link).
 Loc.: Tonga Is.; Fiji Is.; New Caledonia; New Hebrides; North Australia; New Guinea; Solomon Is.; Celebes; Borneo; Malaya; Indonesia; Marshall Is.; Caroline Is.; Mariana Is.; Philippines; Bonin Is. (小笠原); Formosa; Ryukyu; Amami Is.; Andaman Is.; Ceylon; Maldive Is.; Persian Gulf; Red Sea; Mauritius; Madagascar; Zanzibar (East coast of Africa). 奄美以南.
61. *L. (L.) truncata* (Humphrey, 1786) (*Strombus*)
 a. *L. (L.) truncata truncata* (Humphrey, 1786)
 Syn. *Strombus bryonia* Gmelin 1791; *Lambis davilae* Röding, 1798; *L. radix* Röding, 1798; *Pterocera truncata* Lamarck, 1822; *Pyrula bengalina* Grateloup, 1840; *Pterocera (Heptadactylus) radixbryoniae* Mörch, 1852.
 Loc.: Cocos Atolls; Andaman Is.; Ceylon; Seychelles Is.; Mauritius; Madagascar; Zanzibar (East coast of Africa).
 b. *L. (L.) truncata sebae* (Kiener, 1843) (*Pterocera*) ラクダガイ
 Syn. *Pterocera (Heptadactylus) sowerbyi* Mörch, 1872; *Lambis bryonia* Hirase & Pilsbry, 1935 (non Gmelin), *L. truncata* Hirase, 1938 (non Solander); *L. truncata* Butot, 1955 (non Humphrey); *L. truncata* Oyama, 1958 (non Humphrey).
 Loc.: Society Is.; Samoa Is.; Line Is.; New Caledonia; Fiji Is.; Gilbert Is.; North-east Australia; Borneo; Java; Marshall Is.; Mariana Is.; Caroline Is.; Philippines; Ryukyu; Amami Is.; Red Sea. 奄美以南.
62. *L. (L.) crocata* (Link, 1807) (*Pterocera*)
 a. *L. (L.) crocata crocata* (Link, 1807) サソリガイ
 Syn. *Lambis scorpius* Röding, 1798 (non Linné) *Strombus aculeatus* Perry,

1811; *Pterocera aurantia* Lamarck, 1822; *St. lambis* Wood, 1825 (non Linné); *Pteroceras aurantiacum* Sowerby, 1825.

Loc.: Samoa Is.; New Hebrides; New Guinea; Indonesia; Marshall Is.; Caroline Is.; Philippines; Formosa; Ryukyu; Amami Is.; Ceylon; Chagos Is.; Seychelles Is.; Mauritius; Madagascar; Zanzibar (East coast of Africa). 奄美以南.

b. *L. (L.) crocata pilsbryi* Abbott, 1960

Syn. *Pteroceras lambis* Sowerby, 1842 (non Linné, in part)

Loc.: Marquesas Is.

Subgenus *Millepes* Mörch 1852 フシデサソリガイ亜属

Syn. *Millipes* H. & A. Adams 1854, *Millipes* Mörch 1870.

63. *L. (M.) millepeda* (Linné, 1758) (*Strombus*) ムカデガイ

Loc.: New Guinea; Indonesia; Philippines.

64. *L. (M.) digitata* (Perry, 1811) (*Strombus*) ホソムカデガイ

Syn. *Pterocera millepeda* Lamarck, 1816 (non Linné); *Pt. elongata* Swainson, 1821; *Pt. crocea* Reeve, 1842; *Pt. novemdactylis* Deshayes, 1843; *Lambis violacea* Abbott, 1955 (non Swainson).

Loc.: Samoa Is.; Philippines; Mauritius; Madagascar; Mozambique (East coast of Africa).

65. *L. (M.) scorpius* (Linné, 1758) (*Strombus*)

a. *L. (M.) scorpius scorpius* (Linné, 1758) フシデサソリガイ

Syn. *Lambis chiragra* Röding, 1798 (non Linné, non Gmelin), *Pterocera nodosa* Lamarck, 1816.

Loc.: Samoa Is.; Fiji Is.; New Caledonia; Solomon Is.; North Australia; New Guinea; Indonesia; Borneo; Caroline Is.; Philippines; Formosa; Ryukyu; Amami Is. 奄美以南.

b. *L. (M.) scorpius indomaris* Abbott, 1960

Syn. *Strombus sinuatus* Perry, 1811 (non Solander); *Pteroceras pseudoscorpio* Schubert & Wagner, 1829 (non Lamarck).

Loc.: Andaman Is.; Ceylon; Chagos Is.; Mauritius; Seychelles Is.; Madagascar; East coast of Africa.

66. *L. (M.) robusta* (Swainson, 1821) (*Strombus*)

Syn. *Pterocera pseudoscorpio* Lamarck, 1822.

Loc.: Society Is.; Line Is.

67. *L. (M.) violacea* (Swainson, 1821) (*Strombus*) ムラサキムカデガイ

Syn. *Strombus purpureus* Mawe, 1823; *Pterocera purpurea* Swainson, 1825; *Pt. multipes* Sowerby, 1842; *Pt. multipes* Deshayes, 1843.

Loc.: Indonesia; Madagascar; Seychelles; Mauritius; Zanzibar (East coast of Africa).

Subgenus *Harpago* Mörch, 1852 スイジガイ亜属

Syn. *Pterocera* Anton 1839.

68. *L. (H.) chiragra* (Linné, 1758) (*Strombus*)

a. *L. (H.) chiragra chiragra* (Linné, 1758) スイジガイ

Syn. *Lambis harpago* Röding, 1798; *L. undulata* Röding, 1798; *Pterocera rugosum* Sowerby, 1851 (シワクチガイ) *Pt. kochii* Freyer, 1855; *Lambis arthritica* Abbott, 1949 (non Röding).

Loc.: Society Is.; Line Is.; Samoa Is.; New Caledonia; New Hebrides; Solomon Is.; North Australia; New Guinea; Indonesia; Borneo; Malaya; Marshall Is.; Mariana Is.; Caroline Is.; Philippines; Formosa; Ryukyu; Amami Is.; Thailand; Andaman Is.; Ceylon; Cocos Atolls. 奄美以南.

b. *L. (H.) chiragra arthritica* [Röding, 1798] アフリカスイジガイ

Syn. *Strombus divergens* Perry, 1811; *St. nigricans* Perry, 1811; *Pterocera rugosum* Sowerby, 1842 (in part); *Pt. chiragra* Freyer, 1855 (non Linné); *Pt. chiragra* Chenu, 1959 (non Linné); *Pt. (Harpago) chiragra* Thiele, 1929 (non Linné).

Loc.: Maldive Is.; Seychelles Is.; Mauritius; Réunion; Madagascar; East coast of Africa.

Family Aporrhaidae モミジソデガイ科

Genus *Aporrhais* Da Costa 1778 モミジソデガイ属

Syn. *Chenopus* Philippi 1836.

Subgenus *Aporrhais* s. s. モミジソデガイ亜属

*69. *A. (A.) pespelicani* (Linné, 1758) (*Strombus*) モミジソデガイ

Syn. *Rostellaria pespelecani* Lamarck, 1822; *Aporrhais quadrifidus* Da Costa, 1778; *Cirropteron semilunare* Sars, 1835.

Loc.: Europe; Mediterranean Sea to North Norway and Iceland.

*70. *A. (A.) serresianus* (Michaud, 1828) (*Rostellaria*) ヒメモミジボラ

Syn. *Aporrhais pescarbonis* Brogniart, *A. macandear* Jeffreys 1867.

Loc.: Europe; Mediterranean Sea to Norway.

*71. *A. (A.) senegalensis* Gray, 1838

Loc.: Senegal.

Subgenus *Arrhoges* Gabb 1868 アメリカモミジボラ亜属

- *72. *A. (A.) occidentalis* (Beck, 1836) (*Rostellaria*) アメリカモミジボラ
Syn. *Aporrhais occidentalis mainensis* Johnson, 1930; *A. occidentalis labrodorensis* Johnson, 1930.
Loc.: East coast of America.

Family Struthiolariidae ダチヨウノアシボラ科

Genus *Struthiolaria* Lamarck 1812 ダチヨウノアシボラ属

Subgenus *Struthiolaria* s. s. ダチヨウノアシボラ亜属

- *73. *S. papulosa* (Martyn, 1784) (*Buccinum*) ダチヨウノアシボラ (Op. 479)
Ostrich foot shell
Syn. *Struthiolaria nodulosa* Lamarck, 1812; *Struthiolaria gigas* Sowerby, 1842
Loc.: Australia; New Zealand.

Genus *Pellicaria* Gray 1857 ヒメダチヨウノアシボラ属

- *74. *P. vermis* (Martyn, 1784) (*Buccinum*) ヒメダチヨウノアシボラ (Op. 479)
Loc.: New Zealand.
*75. *P. tricarinata* (Lesson, 1830) (*Struthiolaria*)
Loc.: New Zealand.

Genus *Tylospira* Harris 1897

- *76. *T. scutulata* (Gmelin, 1791) (*Buccinum*)
Loc.: N. S. Wales, Australia.

Genus *Perissodonta* Martens 1878

- *77. *P. mirabilis* (Smith, 1875) (*Struthiolaria*)
Loc.: Kerguelen Is.